

A COMMON FIXED POINT THEOREM FOR SIX SELF MAPPING UNDER S-B PROPERTY

Dr. V. K. Gupta*

Swati Choursiya**

Dr. V. H. Badshah***

Abstract

In this paper we use the S-B property with semi compatibility, weakly compatibility and prove common fixed point theorem for six self mapping on a fuzzy metric space.

Keywords- Weakly compatible, semi compatible, (S-B) property, Fuzzy metric space

Zadeh [15] was the foundation of fuzzy sets in 1965. To use this concept in topology and analysis, many authors have extensively developed the theory of fuzzy sets and its applications. Especially, Deng [2], Erceg [3], and Kramosil and Michalek [8] have introduced the concepts of fuzzy metric spaces in different ways. In 1988, Grabiec [7] extended the fixed point theorem in fuzzy metric space. George and Veeramani [5], [6] have modified the concept of fuzzy metric space introduced by Kramosil and Michalek. They have also shown that every metric induces a fuzzy metric. Singh et al. [14] proved various fixed point theorems using the concepts of semi-compatibility, compatibility in Fuzzy metric space.

In this paper, we proved common fixed point theorem for six self mapping by using the semi compatibility, weakly compatibility and new (S-B) property in fuzzy metric space, and using the result from [9] and [11]

* Govt. Madhav Science PG College, Vikram University, Ujjain (M.P), India

** School of Studies in Mathematics, Vikram University, Ujjain, (M.P), India

*** Prof. and Head, School of Studies in Mathematics, Vikram University, Ujjain, (M.P), India

Preliminaries

Definition1. [15] Let X be any set. A fuzzy set in X is a function with domain X and values in $[0, 1]$.

Definition2. [11] A binary operation $*$: $[0, 1] \times [0, 1] \rightarrow [0, 1]$ is said to be a continuous t -norm if $*$ satisfies the following conditions: For $a, b, c, d \in [0, 1]$,

- (i) $*$ is associative and commutative,
- (ii) $*$ is continuous,
- (iii) $a * 1 = a$, for all $a \in [0, 1]$
- (iv) $a * b \leq c * d$ whenever $a \leq c$ and $b \leq d$, for each $a, b, c, d \in [0, 1]$.

We observe that $a * b = \min \{a, b\}$ is a t -norm.

Definition3. [8] The triplet $(X, M, *)$ is said to be a **Fuzzy metric space** if X is an arbitrary set, $*$ is a continuous t -norm and M is a Fuzzy set on $X \times X \times [0, \infty] \rightarrow [0, 1]$, satisfying the following conditions, for all $x, y, z \in X$ and $s, t > 0$.

- (FM-1) $M(x, y, 0) = 0$,
- (FM-2) $M(x, y, t) = 1$ for all $t > 0$ if and only if $x = y$,
- (FM-3) $M(x, y, t) = M(y, x, t)$,
- (FM-4) $M(x, y, t) * M(y, z, s) \leq M(x, z, t+s)$,
- (FM-5) $M(x, y, \cdot) : [0, \infty] \rightarrow [0, 1]$, is left continuous,
- (FM-6) $\lim_{n \rightarrow \infty} M(x, y, t) = 1$.

Note that can be considered as the degree of nearness between x and y with respect to t . We identify $x = y$ with $M(x, y, t) = 1$ for all $t > 0$.

Definition4. [5] Let $(X, M, *)$ be fuzzy metric space then,

- a) A sequence $\{x_n\}$ in X is said to be **convergent** to x in X if for each $\varepsilon > 0$ and each $t > 0$, there exists $n_0 \in \mathbb{N}$ such that $M(x_n, x, t) > 1 - \varepsilon$ for all $n \geq n_0$.
- b) A sequence $\{x_n\}$ in X is said to be **Cauchy sequence** for each $\varepsilon > 0$ and $t > 0$, there exists $n_0 \in \mathbb{N}$ such that $M(x_m, x_n, t) > 1 - \varepsilon$ for all $m, n \geq n_0$.
- c) A fuzzy metric space in which every Cauchy sequence is convergent is said to be **complete**.

Definition5. [9] Self mappings A and S of a Fuzzy metric space $(X, M, *)$ are said to be **compatible** if and only if $M(ASx_n, SAx_n, t) \rightarrow 1$ for all $t > 0$, whenever $\{x_n\}$ is a sequence in X such that $Sx_n, Ax_n \rightarrow p$ for some p in X as $n \rightarrow \infty$.

Definition6. [14] Two self mapping A and S of a Fuzzy metric space (X, d) are said to be **weakly compatible** if they commute at their coincidence points i.e., if $Ax = Sx$ then $ASx = SAx$.

Definition7. [14] Two self mapping A and S of a Fuzzy metric space (X, d) are said to be **semi-compatible** if $\lim_{n \rightarrow \infty} d(ASx_n, Sx_n, a) = 0$ for all $a \in X$, where $\{x_n\}$ is a sequence in X such that if $\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} Sx_n = x$ for some x in X .

Definition8. ([13], [12]) Let S and T be two self mappings of a fuzzy metric space $(X, M, *)$. We say that S and T satisfy the property $S-B$ if there exists a sequence $\{x_n\}$ in X such that $\lim_{n \rightarrow \infty} Sx_n = \lim_{n \rightarrow \infty} Tx_n = z$ for some $z \in X$.

Lemma1. [4] Let $(X, M, *)$ be a fuzzy metric space. For all $x, y \in X$, $M(x, y, \bullet)$ is non decreasing.

Lemma2. [1] Let $(X, M, *)$ be a fuzzy metric space if there exists $k \in (0, 1)$ such that $M(x, y, kt) \geq M(x, y, t)$ then $x=y$.

Example1. [13] Let $X = [0, +\infty)$. Define $A, B: X \rightarrow X$ by $Ax = x + 1/2$ and $Bx = 2x + 1/2$, $x \in X$. Consider the sequence $x_n = 1/n$, clearly $\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} Bx_n = 1/2$. Then A and B satisfy $(S-B)$ property.

Example2. Let $X = [2, +\infty)$. Define $A, B: X \rightarrow X$ by $Ax = 2x + 1$ and $Bx = x + 1$, for all $x \in X$. Suppose property $(S-B)$ holds, then there exists a sequence $\{x_n\}$ in X satisfying

$$\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} Bx_n = z \text{ for some } z \in X.$$

Therefore $\lim_{n \rightarrow \infty} x_n = z - 1$ and $\lim_{n \rightarrow \infty} x_n = (z - 1)/2$.

Then $z = 1$, which is a contradiction since $1 \notin X$. Hence A and B do not satisfy the property $(S-B)$.

Rajinder Sharma [12] proved the following:

Theorem 1.1 Let $(X, M, *)$ be a fuzzy metric space with $t*t \geq t$ for all $t \in [0,1]$ and condition (FM-6). Let A, B, S and T be mappings of X into itself such that

(1.1.1) $A(X) \subset T(X)$ and $B(X) \subset S(X)$,

(1.1.2) (A, S) or (B, T) satisfies the property $(S-B)$,

(1.1.3) there exists a constant $k \in (0, 1)$ such that

$$M^{2p}(Ax, By, kt) \geq \min\{M^{2p}(Sx, Ty, t), M^q(Sx, Ax, t), M^{q'}(Ty, By, t), M^r(Sx, By, t), M^{r'}(Ty, Ax, (2-\alpha)t), M^s(Sx, Ax, t),$$

$$M^{s'}(Ty, Ax, (2-\alpha)t), M^l(Sx, By, t), M^{l'}(Ty, By, t)\}$$

for all $x \in X$, $\alpha \geq 0, \alpha \in (0, 2)$, $t > 0$ and $0 < p, q, q', r, r', s, s', l, l' \leq 1$ such that

$$2p = q + q' = r + r' = s + s' = l + l'.$$

(1.1.4) the pairs (A, S) and (B, T) are weakly compatible,

(1.1.5) one of (X) , $B(X)$, $S(X)$ or $T(X)$ is a closed subset of X , then A, B, S and T have a unique common fixed point in X .

N. Umadevi [10] proved the following:

Theorem 1.2 Let $(X, M, *)$ be a fuzzy metric with $*$ is min t- norm with condition (1). Let A, B, S and T be mappings of X into itself such that

(1.2.1) $A(X) \subset T(X)$, $B(X) \subset S(X)$, $T(X)$ is closed subset of X ,

(1.2.2) $\{B, T\}$ satisfies the property (S-B),

(1.2.3) there exist a constant $k \in (0, 1)$ and $\alpha \in (0, 2)$ such that $k < \alpha$, $k + \alpha < 2$ and satisfies

$$M(Ax, By, kt) \geq \min\{M(Sx, Ty, t)*M(Sx, Ax, t)*M(Ty, By, t)*M(Sx, By, t)*M(Ty, Ax, (2-\alpha)t)\}$$

for all $t > 0$.

(1.2.4) (A, S) and (B, T) are weakly compatible,

Then A, B, S and T have a unique common fixed point in X .

Main Result

Theorem 1.3 Let $(X, M, *)$ be a fuzzy metric with $a*a \geq a$ for all $a \in [0, 1]$ and the condition (fm6). Let A, B, S, T, P and Q be mappings from X into itself such that

(1.3.1) $P(X) \subset AB(X)$, $Q(X) \subset ST(X)$,

(1.3.2) the pairs (A, B) , (S, T) , (Q, B) , (T, P) and are commuting mappings,

(1.3.3) the pairs (P, ST) are semi-compatible and (Q, AB) are weakly compatible,

(1.3.4) $\{P, ST\}$ or $\{Q, AB\}$ satisfies the property (S-B),

$$(1.3.6) M(Px, Qy, qt) \geq M(STx, ABy, t) * M(Px, STx, t) * M(Qy, ABy, t) * M(Px, ABy, t),$$

(1.3.5) one of $AB(X)$, $P(X)$, $AB(X)$ or $Q(X)$ is closed subset of X ,

Then A, B, S, T, P and Q have a unique common fixed point in X .

Proof: Without loss of generality we suppose that (Q, AB) satisfies the S-B property, so there exists a sequence $\{x_n\}$ in X such that $\lim_{n \rightarrow \infty} Qx_n = \lim_{n \rightarrow \infty} ABx_n = z$ for some $z \in X$.

Since $Q(X) \subset ST(X)$ there exists a sequence $\{y_n\}$ in X such that $Qx_n = STy_n$. So $\lim_{n \rightarrow \infty} STy_n = z$.

Now we prove that $\lim_{n \rightarrow \infty} Py_n = z$.

By (1.3.6) we get,

$$M(Py_n, Qx_n, qt) \geq M(STy_n, ABx_n, t) * M(Py_n, STy_n, t) * M(Qx_n, ABx_n, t) * M(Py_n, ABx_n, t),$$

$$M(Py_n, z, qt) \geq M(z, z, t) * M(Py_n, z, t) * M(z, z, t) * M(Py_n, z, t),$$

Therefore by lemma 2,
$$M(Py_n, z, qt) \geq M(Py_n, z, t)$$

we have $Py_n = z$.

Since (P, ST) are semi compatible pairs, we have $PSTy_n = Pz$ and $STPy_n = STz$.

Since the limit in fuzzy metric space is unique, we get $Pz = STz$.

By (1.3.6) we get,

$$M(PPy_n, Qx_n, qt) \geq M(STPy_n, ABx_n, t) * M(PPy_n, STy_n, t) * M(Qx_n, ABx_n, t) * M(PPy_n, ABx_n, t)$$

Taking limit $n \rightarrow \infty$, we have

$$M(Pz, z, qt) \geq M(Pz, z, t) * M(Pz, z, t) * M(z, z, t) * M(Pz, z, t)$$

Therefore by lemma 2,
$$M(Pz, z, qt) \geq M(Pz, z, t).$$

we have $Pz = STz = z$.

By (1.3.6) we get,

$$M(PTz, Qy_n, qt) \geq M(STTz, ABy_n, t) * M(PTz, STTz, t) * M(Qy_n, ABy_n, t) * M(PTz, ABy_n, t),$$

Since $ST = TS$, $PT = TP$, so we have

$$STTz = (TS)Tz = T(STz) = Tz \text{ and } PTz = T(Pz) = Tz, \text{ we have}$$

Taking limit $n \rightarrow \infty$, we have

$$M(Tz, z, qt) \geq M(Tz, z, t) * M(Tz, Tz, t) * M(z, z, t) * M(Tz, z, t),$$

Therefore by lemma 2,
$$M(Tz, z, qt) \geq M(Tz, z, t)$$

we have $Tz = z$. So we have $Pz = z$.

hence $Tz = Pz = Sz = z$.

Suppose $AB(X)$ is closed subset of X . there exists $u \in X$ such that $z = ABu$.

By (1.3.6) we get,

$$M(Py_n, Qu, qt) \geq M(STy_n, ABu, t) * M(Py_n, STx_n, t) * M(Qu, ABu, t) * M(Py_n, ABu, t),$$

Taking limit $n \rightarrow \infty$, we have

$$M(z, Qu, qt) \geq M(z, z, t) * M(z, z, t) * M(Qu, z, t) * M(z, z, t),$$

Therefore By lemma 2
$$M(z, Qu, qt) \geq M(Qu, z, t),$$

hence $Qu = z = ABu$.

Since (Q, AB) are weakly compatible, we have $QABu = ABQu$. Thus $Qz = ABz$.

By (1.3.6) we get,

$$M(Py_n, Qz, qt) \geq M(STy_n, ABz, t) * M(Py_n, STy_n, t) * M(Qz, ABz, t) * M(Py_n, ABz, t),$$

Taking limit $n \rightarrow \infty$, we have

$$M(z, Qz, qt) \geq M(Qz, Qz, t) * M(z, Qz, t) * M(Qz, Qz, t) * M(z, Qz, t),$$

Therefore By lemma 2
$$M(z, Qz, qt) \geq M(Qz, Qz, t)$$

Thus $Qz = z$.

By (1.3.6) we get,

$$M(Py_n, QBz, qt) \geq M(STy_n, ABBz, t) * M(Py_n, STy_n, t) * M(QBz, ABBz, t) * M(Py_n, ABBz, t),$$

Since $AB = BA$ and $QB = BQ$

$$ABBz = B(ABz) = Bz \text{ and } QBz = BQz = Bz.$$

Taking limit $n \rightarrow \infty$, we have

$$\text{So } M(z, Bz, qt) \geq M(z, Bz, t) * M(z, z, t) * M(Bz, Bz, t) * M(z, Bz, t),$$

Therefore By lemma 2
$$M(z, Bz, qt) \geq M(z, Bz, t)$$

Thus $Bz = z$.

So $Az = z$ thus $Qz = Az = Bz = z$.

Hence z , is the common fixed point of A, B, P, Q, S and T .

Uniqueness Let w is another common fixed point of A, B, P, Q, S and T

By (1.3.6) we get,

$$M(Pz, Qw, qt) \geq M(STz, ABw, t) * M(Pz, STz, t) * M(Qw, ABw, t) * M(Pz, ABw, t),$$

$$M(z, w, qt) \geq M(z, w, t) * M(z, z, t) * M(w, w, t) * M(z, w, t),$$

Therefore By lemma 2
$$M(z, w, qt) \geq M(z, w, t)$$

Hence $z = w$.

References

- [1] Cho, Y.J.: Fixed points in fuzzy metric spaces. *J. Fuzzy Math.* 5(4), 949-962 (1997)
- [2] Deng, Z. K.: Fuzzy pseudo metric spaces, *J. Math. Anal. Appl.*, 86, 74-95 (1982)
- [3] Erceg, M. A.: Metric spaces in fuzzy set theory, *J. Math. Anal. Appl.*, 69, 205-230 (1979)
- [4] Grabiec, M.: Fixed point in fuzzy metric spaces, *Fuzzy Sets and Systems*, 27, 385-389 (1988)
- [5] George, A. and Veeramani, P.: On some results in fuzzy metric spaces, *Fuzzy Sets and Systems*, 64, 395-399 (1994)
- [6] George, A. and Veeramani, P.: On some results of analysis for fuzzy metric spaces, *Fuzzy Sets and Systems*, 90, 365-368 (1997)
- [7] Grabiec, M.: Fixed points in fuzzy metric space. *Fuzzy sets and systems*, 27, 385-389 (1988)
- [8] Kramosil, I. and Michalek, J.: Fuzzy metric and statistical metric spaces. *Ky-bernetika.* 11, 336-344 (1975)
- [9] Mishra, S. N., Sharma, N. and Singh, S. L.: Common fixed points of maps in fuzzy metric spaces. *Internat. J. Math. Math. Sci.*, 17, 253-258 (1994)
- [10] Sastry, K. P. R., Naidu, G. A. and Umadevi, N.: A common fixed point theorem for four self maps on a fuzzy metric space under S-B property, *International Journal of Mathematical Archive.* 4(1), 271-275 (2013)
- [11] Schweizer, B. and Sklar. A.: *Probabilistic Metric Spaces.* North Holland, Amsterdam (1983)
- [12] Sharma, R.: Common fixed point of weakly compatible mappings under a new property in fuzzy metric spaces. *Network and Complex Systems.* vol2, no4, 38 – 44 (2012)
- [13] Sharma, Sushil and Bamboria, D.: Some new common fixed point theorems in fuzzy metric space under strict contractive conditions. *J. Fuzzy Math.* 14, No.2, 1-11, (2006)
- [14] Singh, B. and Jain, S.: Weak compatibility and fixed point theorems in fuzzy metric spaces. *Ganita.* 56(2), 167-176 (2005)
- [15] Zadeh, L.A.: Fuzzy sets. *Infor. And Control.* 8, 338-353 (1965)